

**International School of Economics at TSU**  
**Econometrics II**  
**Lasha Chochua**

**Problem Set 1**

**Instructions:** You are encouraged to solve the problems before the recitation. Additionally, you are encouraged to work in groups. It is **not mandatory** to submit solutions unless stated otherwise. However, if you would like to share your solution, I would be happy to review it.

**Problem 1:** Consider the stacked system of linear equations written in matrix notation as:

$$\mathbf{Y} = \bar{\mathbf{X}}\beta + \mathbf{e},$$

where:

- $\mathbf{Y} \in \mathbb{R}^{nm \times 1}$  is the stacked outcome vector,
- $\bar{\mathbf{X}} \in \mathbb{R}^{nm \times \bar{k}}$  is the block-diagonal regressor matrix,
- $\beta \in \mathbb{R}^{\bar{k} \times 1}$  is the parameter vector,
- $\mathbf{e} \in \mathbb{R}^{nm \times 1}$  is the stacked error vector.

Assume the following:

- $\mathbb{E}[\mathbf{e} | X] = 0$
- $\mathbb{E}[\mathbf{e}\mathbf{e}'] = \Omega = I_n \otimes \Sigma$ , with  $\Sigma \in \mathbb{R}^{m \times m}$  a symmetric positive definite matrix.

Derive the Generalized Least Squares (GLS) estimator for  $\beta$ , and show that it can be written as:

$$\hat{\beta}_{\text{glS}} = \left( \bar{\mathbf{X}}' (I_n \otimes \Sigma^{-1}) \bar{\mathbf{X}} \right)^{-1} \left( \bar{\mathbf{X}}' (I_n \otimes \Sigma^{-1}) \mathbf{Y} \right).$$

**Problem 2:** Consider the stacked system

$$\mathbf{Y} = \bar{\mathbf{X}}\beta + \mathbf{e},$$

where

- $\mathbf{Y}$  is the  $nm \times 1$  stacked vector of dependent variables,
- $\bar{\mathbf{X}}$  is the  $nm \times \bar{k}$  stacked regressor matrix,

- $\mathbf{e}$  is the  $nm \times 1$  stacked error vector.

Assume

$$E[\mathbf{e} \mid X] = 0$$

and

$$E[\mathbf{e}\mathbf{e}' \mid X] = I_n \otimes \Sigma,$$

where  $\Sigma$  is an  $m \times m$  positive definite matrix.

The GLS estimator is

$$\hat{\beta}_{gls} = \left( \sum_{i=1}^n \bar{X}_i' \Sigma^{-1} \bar{X}_i \right)^{-1} \left( \sum_{i=1}^n \bar{X}_i' \Sigma^{-1} Y_i \right).$$

Show that

$$\text{Var}(\hat{\beta}_{gls} \mid X) = \left( \sum_{i=1}^n \bar{X}_i' \Sigma^{-1} \bar{X}_i \right)^{-1}.$$

**Problem 3:** Suppose the covariance matrix  $\Sigma$  in the system model is unknown. We consider the same system as in Problem 1:

$$\mathbf{Y} = \bar{\mathbf{X}}\beta + \mathbf{e},$$

We still assume:

- $\mathbb{E}[\mathbf{e} \mid X] = 0$
- $\mathbb{E}[\mathbf{e}\mathbf{e}' \mid X] = \Omega = I_n \otimes \Sigma$
- $\Sigma$  is **unknown**, but we have a **consistent estimator**  $\hat{\Sigma}$

1. Write the **feasible GLS (FGLS)** estimator,  $\hat{\beta}_{\text{sur}}$ , using matrix notation.
2. Prove that under regularity conditions (e.g. Assumption 7.2 and consistency of  $\hat{\Sigma}$ ), the FGLS estimator is **asymptotically normal**:

$$\sqrt{n}(\hat{\beta}_{\text{sur}} - \beta) \xrightarrow{d} \mathcal{N}(0, V_{\beta}^*)$$

where

$$V_{\beta}^* = \left( \text{plim} \frac{1}{n} \bar{\mathbf{X}}' (I_n \otimes \Sigma^{-1}) \bar{\mathbf{X}} \right)^{-1}.$$

**Problem 4:** Use the data in NBASAL.RAW to answer this question.

- a. Estimate an SUR model for the three response variables points, rebounds, and assists. The explanatory variables in each equation should be age, exper, exper2, guard, forward, black, and marr. Does marital status have a positive or negative effect on each variable? Is it statistically significant in the assists equation?
- b. Test the hypothesis that marital status can be excluded entirely from the system.